

FEZ1 Rabbit pAb

货号: **AYP18481**

产品信息

| | |
|-------|--|
| 反应 | Human |
| 宿主 | Rabbit |
| 克隆性 | Polyclonal |
| 预测反应 | |
| 应用 | WB |
| 推荐浓度 | WB: 1:200 - 1:2000 |
| 理论分子量 | 12kDa/45kDa |
| 实测分子量 | 71kDa |
| 形式 | Liquid |
| 保存条件 | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal, 50% glycerol, pH7.3. |
| 偶联物 | Unconjugated |
| 阳性对照 | U-251MG |
| 细胞定位 | Cell membrane, Cytoplasm, centrosome, cytoskeleton, microtubule organizing center |
| 纯化 | Affinity purification |

抗原信息


| | |
|------|---|
| 抗原信息 | Recombinant fusion protein containing a sequence corresponding to amino acids 1-160 of human FEZ1 (NP_005094.1). |
| 序列 | MEAPLVSLDEEFEDLRPSCSEDPEEKPQCFYGSSPHHLEDPSLSELENFSSEIISFKSMEDLVNEFDEKLNVCFRNYNAKTE NLAPVKNQLQIQEEEEETLQDEEVWDALTDNYIPSLSEDWRDPNIEALNGNCSDTEIHEKEEEEFNEKSENDSGINEEP |

靶点信息

| | |
|------|--|
| 研究背景 | This gene is an ortholog of the <i>C. elegans</i> unc-76 gene, which is necessary for normal axonal bundling and elongation within axon bundles. Expression of this gene in <i>C. elegans</i> unc-76 mutants can restore to the mutants partial locomotion and axonal fasciculation, suggesting that it also functions in axonal outgrowth. The N-terminal half of the gene product is highly acidic. Alternatively spliced transcript variants encoding different isoforms of this protein have been described. |
| 基因ID | 9638 |

| | |
|-------|-------------|
| 基因名 | FEZ1 |
| Swiss | Q99689 |
| 别名 | FEZ1;UNC-76 |

产品验证



Western blot analysis of FEZ1 expressed in U-251MG using FEZ1 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn