

# PDCD2 Rabbit pAb

货号: AYP18529

## 产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	<b>WB:</b> 1:200 - 1:2000
理论分子量	21kDa/22kDa/25kDa/35kDa/38kDa
实测分子量	38kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal, 50% glycerol, pH7.3.
偶联物	Unconjugated
阳性对照	293T, HeLa
细胞定位	Nucleus
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-344 of human PDCD2 (NP_002589.2).
序列	MAAAGARPVELGFAESAPAWRLRSEQFSPKVGGRPAWLGAAGLPGPQALACELCGRPLSFLQVYAPLPGRPDFAHRCIF LFCCREQPCCAGLRVFRNQLPRKNDYFSEPPSENPPPETGESVCLQLKSGAHLRCVCGCLGPKTCSRCHKAYYCSKEHQ LDWRLGHKQACAQPDHLDHIIPDHNFLPFEFEIVIEDEIMPEVVEKEDYSEIIGSMGEALEEELDSMAKHESREDKIFQKF KTQIALEPEQILRYGRGIAPWISGENIPQEKDIPDCPGAKRILEFQVMPQLLNLYLKADRLGKSIDWGLAVFTCAESCSLGT GYTEEFVWKQDVTDTTP

## 靶点信息

研究背景	This gene encodes a nuclear protein expressed in a variety of tissues. Expression of this gene has been shown to be repressed by B-cell CLL/lymphoma 6 (BCL6), a transcriptional repressor required for lymph node germinal center development, suggesting that BCL6 regulates apoptosis by its effects on this protein. Alternative splicing results in multiple transcript variants and pseudogenes have been identified on chromosomes 9 and 12.
------	---

基因ID	5134
基因名	PDCD2
Swiss	Q16342
别名	PDCD2;RP8;ZMYND7

### 产品验证



Western blot analysis of PDCD2 expressed in 293T, HeLa using PDCD2 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

### 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)