

IK Rabbit pAb

货号: **AYP18537**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:200 - 1:2000
理论分子量	65kDa
实测分子量	75kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	A-549,HeLa,LO2,Mouse brain,Mouse testis,Rat kidney
细胞定位	Nucleus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-280 of human IK (NP_006074.2).
序列	MPERDSEPFNSNPLAPDGHVDVDDPHSFHQSKLTNEDFRKLLMTPRAAPTSAPPSKSRHHEMPREYNEDEDPAARRRKKKS YYAKLRQQEIERERELAEKYRDRAKERRDGVNKDYEETELISTTANYRAVGPTAEADKSAEKRRQLIQESKFLGGDMEHT HLVKGLDFALLQKVRAEIASKEKEEELMEKPQKTKKDEDPENKIEFKTRLGRNVYRMLFKSKAYERNELFLPGRMAYVVD LDDEYADTDIPTTLIRSKADCPTMEAQTTLTTNDIVIS

靶点信息

研究背景	The protein encoded by this gene was identified by its RED repeat, a stretch of repeated arginine, glutamic acid and aspartic acid residues. The protein localizes to discrete dots within the nucleus, excluding the nucleolus. Its function is unknown. This gene maps to chromosome 5; however, a pseudogene may exist on chromosome 2.
------	--

基因ID	3550
基因名	IK
Swiss	Q13123
别名	IK;CSA2;RED;RER

产品验证



Western blot analysis of IK expressed in Mouse brain,Rat kidney,A-549 using IK Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn