

BTF3 Rabbit pAb

货号: **AYP18554**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:1000
理论分子量	17kDa/22kDa
实测分子量	18KDa/22KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293F,NIH/3T3,Rat testis,PC-12
细胞定位	Cytoplasm,Nucleus
纯化	Affinity purification

抗原信息


抗原信息	A synthetic peptide corresponding to a sequence within amino acids 50-150 of human BTF3 (NP_001198.2).
序列	KLGVNNISGIEEVNMFTNQGTVIHFNNPKVQASLAANTFTITGHAETKQLTEMLPSILNQLGADSLTSLRRLAEALPKQSVD GKAPLATGEDDDDEVPDLV

靶点信息

研究背景	This gene encodes the basic transcription factor 3. This protein forms a stable complex with RNA polymerase IIB and is required for transcriptional initiation. Alternative splicing results in multiple transcript variants encoding different isoforms. This gene has multiple pseudogenes.
基因ID	689

基因名	BTF3
Swiss	P20290
别名	BTF3;BETA-NAC;BTF3a;BTF3b;NACB

产品验证



Western blot analysis of BTF3 expressed in Rat testis,293F,NIH/3T3 using BTF3 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn