

EIF5B Rabbit pAb

货号: **AYP18626**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200
理论分子量	138kDa
实测分子量	179kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	DU145,HeLa,Jurkat,LO2
细胞定位	Cytoplasm
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-273 of human EIF5B (NP_056988.3).
序列	MGKKQKNKSEDSTKDDIDLALAAEIEGAGAAKEQEPQKSKGKKKKEKKKQDFDEDDILKELEELSLEAQGIKADRETVA VKPTENNEEEFTSKDKKKKGQKGGKQSFDDNDSELEDKDSKSKKTAKPKVEMYSGSDDDDDFNKLPKKAKGKAQKSN KKWDGSEEDEDNSKKIKERSRINSSGESGDESDEFQSRKGQKKNQKNKPGPNIESGNEDDDASFKIKTVAQKKAEEKK RERKKRDEEKAKLRKLKEKEELETGKKDQSKQKESQ

靶点信息

研究背景	Accurate initiation of translation in eukaryotes is complex and requires many factors, some of which are composed of multiple subunits. The process is simpler in prokaryotes which have only three initiation factors (IF1, IF2, IF3). Two of these factors are conserved in eukaryotes: the homolog of IF1 is eIF1A and the homolog of IF2 is eIF5B. This gene encodes eIF5B. Factors eIF1A and eIF5B interact on the ribosome along with other initiation factors and GTP to position the initiation methionine tRNA on the start codon of the mRNA so that translation initiates accurately.
基因ID	9669
基因名	EIF5B
Swiss	O60841
别名	EIF5B;IF2

产品验证



Western blot analysis of EIF5B expressed in DU145, HeLa, Jurkat using EIF5B Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn