

BRE Rabbit pAb

货号: **AYP18627**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:100 IF/ICC: 1:50 - 1:200
理论分子量	42kDa/43kDa/46kDa
实测分子量	44kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,HT-29,mouse brain,mouse heart,rat testis
细胞定位	Cytoplasm,Nucleus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 51-229 of human BRE (NP_954661.1).
序列	PNCDRFKLHIPPYAGETLKWDIIFNAQYPELPPDFIFGEDAEFLPDPSALQNLASWNPSPNECLLLVVKELVQQYHQFQCSRL RESSRLMFYQTLLEEPQYGENMEIYAGKKNWWTGEFSARFLLKLPVDFSNIPTYLLKDVNEDPGEDVALLSVSFEDTEAT QVYPKLYLSPRIEHAL

靶点信息

研究背景	This gene encodes an anti-apoptotic, death receptor-associated protein that interacts with tumor necrosis factor-receptor-1. The encoded protein acts as an adapter in several protein complexes, including the BRCA1-A complex and the BRISC complex. The BRCA1-A complex possesses ubiquitinase activity and targets sites of double strand DNA breaks, while the BRISC complex exhibits deubiquitinase activity and is involved in mitotic spindle assembly. This gene is upregulated in several types of cancer.
------	--

基因ID	9577
基因名	BABAM2
Swiss	Q9NXR7
别名	BRE;BRCC4;BRCC45

产品验证



Western blot analysis of BRE expressed in mouse brain, rat testis, HeLa using BRE Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn