

NFIC Rabbit pAb

货号: **AYP18653**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	45kDa/47kDa/48kDa/54kDa/55kDa
实测分子量	60kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,LO2,MCF-7,DU145,NIH/3T3,mouse liver,mouse skeletal muscle,rat liver
细胞定位	Nucleus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-220 of human NFIC (N P_005588.2).
序列	MYSSPLCLTQDEFHPFIEALLPHVRAFAYTWFNLQARKRKYFKKHEKRMSKDEERAVKDELLGEKPEVKQKWASRLAKLR KDIRPECREDFVLSITGKKAPGCVLSNPDQKGMRRIDCLRQADKVVRLDLVMVILFKGIPLLESTDGERLVKAAQC GHPVL CVQPHHIGVAVKELDLYLAYFVRERDAEQSGSPRTGMGSDQEDSKPITLDTTDFQESF

靶点信息

研究背景	The protein encoded by this gene belongs to the CTF/NF-I family. These are dimeric DNA-binding proteins, and function as cellular transcription factors and as replication factors for adenovirus DNA replication. Alternatively spliced transcript variants encoding different isoforms have been described for this gene.
基因ID	4782

基因名	NFIC
Swiss	P08651
别名	NFIC;CTF;CTF5;NF-I;NFI

产品验证



Western blot analysis of NFIC expressed in mouse liver, rat liver, HeLa using NFIC Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn