

UNC5B Rabbit pAb

货号: **AYP18694**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:100
理论分子量	102kDa/103kDa
实测分子量	104kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	U-87MG,A-549,293T,Mouse brain,Mouse lung,Mouse kidney,Rat brain,Rat lung
细胞定位	Cell membrane,Membrane raft,Single-pass type I membrane protein
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 290-377 of human UNC 5B (NP_734465.2).
序列	FQKTACTTICPVDGAWTEWSKWSACSTECAHWRSRECMAPPQNGGRDCSGTLLDSKNCTDGLCMQNKKTLSDPNSH LLEASGDAALY

靶点信息

研究背景	This gene encodes a member of the netrin family of receptors. This particular protein mediates the repulsive effect of netrin-1 and is a vascular netrin receptor. This encoded protein is also in a group of proteins called dependence receptors (DpRs) which are involved in pro- and anti-apoptotic processes. Many DpRs are involved in embryogenesis and in cancer progression. Two alternatively spliced transcript variants have been described for this gene.
------	--

基因ID	219699
基因名	UNC5B
Swiss	Q8IZJ1
别名	UNC5B;UNC5H2;p53RDL1

产品验证



Western blot analysis of UNC5B expressed in Mouse brain,Rat brain,U-87MG using UNC5B Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn