

OCIAD2 Rabbit pAb

货号: **AYP18695**

产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200
理论分子量	11kDa/16kDa
实测分子量	17kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal, 50% glycerol, pH7.3.
偶联物	Unconjugated
阳性对照	293T, LO2, A-549
细胞定位	Endosome
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-99 of human OCIAD2 (NP_689611.1).
序列	MASASARGNQDKDAHFPSPKQSLFLCPKSLHIIHRAEISKIMRECQEESFWKRALPFSLVSMMLVTQGLVYQGYLAANSRF GSLPKVARTASLPVRNAK

靶点信息

研究背景	Predicted to be involved in endocytosis; hematopoietic stem cell homeostasis; and positive regulation of receptor signaling pathway via JAK-STAT. Predicted to act upstream of or within response to bacterium. Predicted to be located in Golgi apparatus; endosome; and lysosome.
基因ID	132299

基因名	OCIAD2
Swiss	Q56VL3
别名	OCIAD2

产品验证



Western blot analysis of OCIAD2 expressed in 293T, LO2, A-549 using OCIAD2 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn