

DDX59 Rabbit pAb

货号: **AYP18713**

产品信息

反应	Human,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200
理论分子量	64kDa/68kDa
实测分子量	64kDa/68kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	
细胞定位	Cytoplasm,Nucleus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-250 of human DDX59 (NP_001026895.2).
序列	MFVPRSLKIKRNANDDGKSCVAKIIKDPEDLQLDKSRDVPVDAVATEAATIDRHISESCFPSPGGQLAEVHSVSPSEQGA KDSHPSEEPVKSFSKTQRWAEPGEPICVCGRYGEYICDKTDEDVCSLECKAKHLLQVKEKEEKSLSNPQKADSEPEP NASYVYKEHPFILNLQEDQIENLKQQLGILVQGQEVTRPIIDFEHCSLPEVLNHNHLLKSGYEVPTPIQMIPVGLLGRDILAS ADTG

靶点信息

研究背景	Predicted to enable RNA binding activity and RNA helicase activity. Predicted to be located in cytoplasm and nucleus. Predicted to be integral component of membrane. Implicated in orofacioidigital syndrome V.
基因ID	83479

基因名	DDX59
Swiss	Q5T1V6
别名	DDX59;OFD5;ZNHIT5

产品验证



Western blot analysis of DDX59 expressed in human liver cancer, rat muscle using DDX59 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn