

— ABLYBIO, Help Your Research



# NTPCR Rabbit pAb

货号: **AYP18716**

## 产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IF/ICC
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IF/ICC:</b> 1:50 - 1:200
理论分子量	20kDa
实测分子量	21kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	U-251MG,A-431,293T
细胞定位	
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 91-190 of human NTPCR (NP_115700.1).
------	--

## 靶点信息

研究背景	The protein encoded by this gene is a non-specific nucleoside triphosphatase that is slow-acting in vitro. This gene is overexpressed in many tumor tissues, and while it is not essential for the cell, overexpression is cytotoxic. However, the cytotoxicity is not related to its triphosphatase activity.
基因ID	84284
基因名	NTPCR
Swiss	Q9BSD7 ( <a href="https://www.uniprot.org/uniprotkb/Q9BSD7/entry">https://www.uniprot.org/uniprotkb/Q9BSD7/entry</a> )
别名	NTPCR,C1orf57,HCR-NTPase,NTPCR Rabbit pAb,Nucleoside triphosphate phosphohydrolase

## 产品验证



Western blot analysis of NTPCR expressed in U-251MG,A-431,293T using NTPCR Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn) (<https://www.ablybio.cn/>[www.ablybio.cn](http://www.ablybio.cn))