

# LRFN3 Rabbit pAb

货号: **AYP18717**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	<b>WB:</b> 1:500 - 1:2000
理论分子量	66kDa
实测分子量	66kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HT-29,Mouse testis,Mouse small intestine,Rat brain
细胞定位	Cell junction,Cell membrane,Cell projection,Single-pass type I membrane protein,axon,dendrite,postsynaptic cell membrane,presynaptic cell membrane,synapse
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 380-530 of human LRFN3 (NP_078785.1).
序列	ELTVGPPPPQLANSTSCDPPRDGDPDALTPPSAASASAKVADTGPPTDRGVQVTEHGATAALVQWPDQRPIPIGIRMYQIQYNSSADDILVYRMIPAESRSFLLDLASGRTYDLCVLAVYEDSATGLTATRPVGCARFSTEPALRPCGAP

## 靶点信息

研究背景	Cell adhesion molecule that mediates homophilic cell-cell adhesion in a Ca <sup>2+</sup> -independent manner. Promotes neurite outgrowth in hippocampal neurons (By similarity).
基因ID	79414

基因名	LRFN3
Swiss	Q9BTN0
别名	LRFN3;FIGLER1;SALM4

## 产品验证



Western blot analysis of LRFN3 expressed in Mouse testis,Rat brain,HT-29 using LRFN3 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)