

FBXO42 Rabbit pAb

货号: **AYP18735**

产品信息

| | |
|-------|--|
| 反应 | Mouse |
| 宿主 | Rabbit |
| 克隆性 | Polyclonal |
| 预测反应 | |
| 应用 | WB |
| 推荐浓度 | WB: 1:500 - 1:2000 |
| 理论分子量 | 77kDa |
| 实测分子量 | 78kDa |
| 形式 | Liquid |
| 保存条件 | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal, 50% glycerol, pH7.3. |
| 偶联物 | Unconjugated |
| 阳性对照 | Mouse lung |
| 细胞定位 | |
| 纯化 | Affinity purification |

抗原信息


| | |
|------|---|
| 抗原信息 | Recombinant fusion protein containing a sequence corresponding to amino acids 468-717 of human FBXO42 (NP_061867.1). |
| 序列 | PSAPEGYDLKIGLSLAPRRGSLPDQKDLRLGSLNWDLKPASSNPMDGMDNRTVGGSMRHPPEQTNGVHTPPHVAS ALAGAVSPGALRRSLEAIKAMSSKGPSASAALSPLGSSPGSPGSQLSSGETVPIPRPGPAQGDGHSPLPIARRLGHHPP QSLNVGKPLYQSMNCKPMQMYVLDIKDTKEKGRVKWVFNSSSVVGPPELHTVVQGRGELIIFGGLMDKKQNVKYY KTNALYFVRAKR |

靶点信息

| | |
|------|--|
| 研究背景 | Members of the F-box protein family, such as FBXO42, are characterized by an approximately 40-amino acid F-box motif. SCF complexes, formed by SKP1 (SKP1A; MIM 601434), cullin (see CUL1; MIM 603134), and F-box proteins, act as protein-ubiquitin ligases. F-box proteins interact with SKP1 through the F box, and they interact with ubiquitination targets through other protein interaction domains (Jin et al., 2004 [PubMed 15520277]).[supplied by OMIM, Dec 2010] |
|------|--|

| | |
|-------|------------------|
| 基因ID | 54455 |
| 基因名 | FBXO42 |
| Swiss | Q6P3S6 |
| 别名 | FBXO42;Fbx42;JFK |

产品验证



Western blot analysis of FBXO42 expressed in Mouse lung using FBXO42 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn