

# DKK2 Rabbit pAb

货号: **AYP18754**

## 产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IF/ICC
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IF/ICC:</b> 1:100 - 1:500
理论分子量	28kDa
实测分子量	30kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal, 50% glycerol, pH7.3.
偶联物	Unconjugated
阳性对照	A-549, NCI-H460
细胞定位	Secreted
纯化	Affinity purification

## 抗原信息


抗原信息	A synthetic peptide corresponding to a sequence within amino acids 100 to the C-terminus of human DKK 2 (NP_055236.1).
序列	CMVCRKKKRCHRDGMCCPSTRCNNGICIPVTESILTPHIPALDGTTRHRDRNHGHYSNHDLGWQNLGRPHTKMSHIKGH EGDPCLRSSDCIEGFCCARHFVTKICKPVLHQGEVCTKQRKKGSHGLEIFQRCDCAKGLSCKVWKDATYSSKARLHVCC KI

## 靶点信息

研究背景	This gene encodes a protein that is a member of the dickkopf family. The secreted protein contains two cysteine rich regions and is involved in embryonic development through its interactions with the Wnt signaling pathway. It can act as either an agonist or antagonist of Wnt/beta-catenin signaling, depending on the cellular context and the presence of the co-factor kremen 2. Activity of this protein is also modulated by binding to the Wnt co-receptor LDL-receptor related protein 6 (LRP6).
------	---

基因ID	27123
基因名	DKK2
Swiss	Q9UBU2
别名	DKK2;DKK-2

## 产品验证



Western blot analysis of DKK2 expressed in A-549, NCI-H460 using DKK2 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)