

TCERG1 Rabbit pAb

货号: **AYP18763**

产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	121kDa/123kDa
实测分子量	160kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal, 50% glycerol, pH7.3.
偶联物	Unconjugated
阳性对照	HeLa, Jurkat, SW480
细胞定位	Nucleus
纯化	Affinity purification

抗原信息


抗原信息	A synthetic peptide corresponding to a sequence within amino acids 550-650 of human TCERG1 (NP_006697.2).
序列	TTRLSMWDRPDDLIGRADVDKIIQEPHKKGMEELKLRHPTPTMLSIQKWQFSMSAIKEEQELMEEINEDEPVKAKKRKR DDNKDIDSEKEAAMEAEIKA

靶点信息

研究背景	This gene encodes a nuclear protein that regulates transcriptional elongation and pre-mRNA splicing. The encoded protein interacts with the hyperphosphorylated C-terminal domain of RNA polymerase II via multiple FF domains, and with the pre-mRNA splicing factor SF1 via a WW domain. Alternative splicing results in multiple transcripts variants encoding different isoforms.
基因ID	10915

基因名	TCERG1
Swiss	O14776
别名	TCERG1;CA150;TAF2S;Urn1

产品验证



Western blot analysis of TCERG1 expressed in HeLa, Jurkat, SW480 using TCERG1 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn