

F8A1 Rabbit pAb

货号: **AYP18789**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	39kDa
实测分子量	39kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	U-251MG,A-549,Mouse spleen,Rat lung,Rat spleen
细胞定位	early endosome,nuclear body,nucleus
纯化	Affinity purification

抗原信息


抗原信息	A synthetic peptide corresponding to a sequence within amino acids 150-250 of human F8A1 (NP_036283.2).
序列	AALRDLGQPAAAAGHFQRAAQLQLPLAALQALGEAASCQLLARDYTGALAVFTRMQRLAREHGSHPVQSLPPPPPPA PQPGPGATPALPAALLPPNSG

靶点信息

研究背景	This gene is contained entirely within intron 22 of the factor VIII gene; spans less than 2 kb, and is transcribed in the direction opposite of factor VIII. A portion of intron 22 (int22h), containing F8A, is repeated twice extragenically closer to the Xq telomere. Although its function is unknown, the observation that this gene is conserved in the mouse implies it has some function. Unlike factor VIII, this gene is transcribed abundantly in a wide variety of cell types.
基因ID	8263

基因名	
Swiss	P23610
别名	F8A1;DXS522E;F8A;HAP40

产品验证



Western blot analysis of F8A1 expressed in Mouse spleen,Rat lung,U-251MG using F8A1 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn