

DYNC1I2 Rabbit pAb

货号: AYP18830

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	68kDa/70kDa/71kDa
实测分子量	71-105kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	SH-SY5Y,U-251MG,Mouse testis,Rat brain
细胞定位	Cytoplasm,cytoskeleton
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 352-612 of human DYN C1I2 (NP_001258717.1).
序列	AAHTHPVYCVNVVGTQNAHNLISITDGGKICSWSLDMLSHPQDSMELVHKQSKAVAVTSMSPVGDVNNFVVGSEEGS VYTACRHGSKAGISEMFEGHQGPITGIHCHAAVGAVDFSHLFTSSFDWTVKLVTTKNNKPLYSFEDNADYVYDVMWSP THPALFACVDGMGRDLWNLNNDTEVPTASISVEGNPALNRVRWTHSGREIAVGDSEGQIVYDVGEQIAVPRNDEWAR FGRTLAEINANRADAEEEEATRIPA

靶点信息

研究背景	This gene encodes a member of the dynein intermediate chain family. The encoded protein is a non-catalytic component of the cytoplasmic dynein 1 complex, which acts as a retrograde microtubule motor to transport organelles and vesicles. A pseudogene of this gene is located on chromosome 10. Alternative splicing results in multiple transcript variants.
------	---

基因ID	1781
基因名	DYNC1I2
Swiss	Q13409
别名	DYNC1I2;DIC74;DNCI2;IC2

产品验证



Western blot analysis of DYNC1I2 expressed in Mouse testis,Rat brain,SH-SY5Y using DYNC1I2 Rabbit p Ab at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120 S.

实验步骤

访问官网浏览详情: www.ablybio.cn