

# RBM4B Rabbit pAb

货号: **AYP18850**

## 产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	<b>WB:</b> 1:500 - 1:2000
理论分子量	40kDa
实测分子量	38kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal, 50% glycerol, pH7.3.
偶联物	Unconjugated
阳性对照	LO2, 293T, U-87MG, Raji, HeLa
细胞定位	Nucleus, nucleolus
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 140-359 of human RBM4B (NP_113680.1).
序列	RMHVQLSTSRLRTAPGMGDQSGCYRCGKEGHWSKECPVDRTGRVADFTEQYNEQYGAVRTPYTMGYGESMYYNDA YGALDYYKRYRVRSYEAVAAAAAASAYNYAEQTMSHLPQVQSTTVTSHLNSTSVDPYDRHLLPNSGAAATSAAMAAA AATTSSYYGRDRSPLRRAAAMLPTVGEQYGYGPESELSQASAATRNSLYDMARYEREQYVDRARYSAF

## 靶点信息

研究背景	Required for the translational activation of PER1 mRNA in response to circadian clock. Binds directly to the 3'-UTR of the PER1 mRNA (By similarity).
基因ID	83759

基因名	RBM4B
Swiss	Q9BQ04
别名	RBM4B;RBM30;RBM4L;ZCCHC15;ZCCHC21B;ZCRB3B

## 产品验证



Western blot analysis of RBM4B expressed in LO2,293T,U-87MG using RBM4B Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)