

LZTS2 Rabbit pAb

货号: **AYP18883**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	72kDa
实测分子量	73kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	DU145,HT-29,HeLa,U-87MG,MCF7,Mouse brain,Rat kidney,Rat brain
细胞定位	Cytoplasm,centrosome,cytoskeleton,microtubule organizing center
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 470-669 of human LZTS 2 (NP_115805.1).
序列	ELVALRVALREARATLRVSEGRARGLQEAAARELELEACSQELQRHRQAEQLREKAGQLDAEAAGLREPPVPPATADPF LLAESDEAKVQRAAAGVGGSLRAQVERLRVELQRERRRGEEQRDSFEGERLAWQAEKEQVIRYQKQLQHNYIQMYRRNR QLEQELQQLSLELEARELADLGLAEQAPCICLEEITATEI

靶点信息

研究背景	The protein encoded by this gene belongs to the leucine zipper tumor suppressor family of proteins, which function in transcription regulation and cell cycle control. This family member can repress beta-catenin-mediated transcriptional activation and is a negative regulator of the Wnt signaling pathway. It negatively regulates microtubule severing at centrosomes, and is necessary for central spindle formation and cytokinesis completion. It is implicated in cancer, where it may inhibit cell proliferation and decrease susceptibility to tumor development. Alternative splicing of this gene results in multiple transcript variants.
------	---

基因ID	84445
基因名	LZTS2
Swiss	Q9BRK4
别名	LZTS2;LAPSER1

产品验证



Western blot analysis of LZTS2 expressed in Mouse brain,Rat kidney,DU145 using LZTS2 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn