

# CNKSR2 Rabbit pAb

货号: **AYP18891**

## 产品信息

|       |  |
|-------|--|
| 反应    | Human  |
| 宿主    | Rabbit   |
| 克隆性   | Polyclonal   |
| 预测反应  |  |
| 应用    | WB   |
| 推荐浓度  | <b>WB:</b> 1:500 - 1:2000  |
| 理论分子量 | 96kDa/101kDa/113kDa/117kDa   |
| 实测分子量 | 117kDa   |
| 形式    | Liquid   |
| 保存条件  | Store at -20°C. Avoid freeze / thaw cycles.<br>Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3. |
| 偶联物   | Unconjugated   |
| 阳性对照  | U-87MG,Jurkat,HepG2  |
| 细胞定位  | Cytoplasm,Membrane,Peripheral membrane protein   |
| 纯化    | Affinity purification  |

## 抗原信息


|      |  |
|------|--|
| 抗原信息 | Recombinant fusion protein containing a sequence corresponding to amino acids 400-750 of human CNK SR2 (NP_055742.2).  |
| 序列   | RKRFNIVEEDTVLYCYEYEKGRSSSQGRRESTPTYGKLRPISMPVEYNWVG DYEDPNKMKRDSRRENSLLRYMSNEKIAQE EYMFQRNSKKDTGKSKKKGDKSNSPHTYLLPSLQMDALRQDIMGTPVPETTLYHTFQQSSLQHKS KKKKNGPIAGKSK RRISCKDLGRGDCEGWLWKKKDAKSYFSQKWKKYWFVLKDASLYWYINEEDEKAEGFISLPEFKIDRASECRKKYAFKAC HPKIKSFYFAAEHLDDMNRWLNRLNMLTAGYAERERIKQEQDYWSESDKEEADTPSTPKQDSPPPPYDTYPRPPSMSCAS PYVEAKHSRLSSTETSQSQSSHEEFRQEV T |

## 靶点信息

|      |   |
|------|---|
| 研究背景 | This gene encodes a multidomain protein that functions as a scaffold protein to mediate the mitogen-activated protein kinase pathways downstream from Ras. This gene product is induced by vitamin D and inhibits apoptosis in certain cancer cells. It may also play a role in ternary complex assembly of synaptic proteins at the postsynaptic membrane and coupling of signal transduction to membrane/cytoskeletal remodeling. Multiple transcript variants encoding different isoforms have been found for this gene. |
|------|---|

|       |                         |
|-------|-------------------------|
| 基因ID  | 22866                   |
| 基因名   | CNKSR2                  |
| Swiss | Q8WXI2                  |
| 别名    | CNKSR2;CNK2;KSR2;MAGUIN |

### 产品验证



Western blot analysis of CNKSR2 expressed in U-87MG, Jurkat, HepG2 using CNKSR2 Rabbit pAb at 1:10 00. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. B locking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

### 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)