

# MYCT1 Rabbit pAb

货号: **AYP18913**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IF/ICC
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IF/ICC:</b> 1:50 - 1:200
理论分子量	26kDa
实测分子量	27kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Rat spleen,Rat lung
细胞定位	Nucleus
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 99-235 of human MYCT 1 (NP_079383.2).
序列	ASAPISQWSSRRSRSSYTHGLNRTGFYRHSGCERRSNLSLASLTFQRQASLEQANSFPRKSSFRSTFHPFLQCPLPVE TESQLVTLPSSNISPTISTSHLSLRPDYWSSNSLRVGLSTPPPPAYESIIFKAFDPS

## 靶点信息

研究背景	May regulate certain MYC target genes, MYC seems to be a direct upstream transcriptional activator. Does not seem to significantly affect growth cell capacity. Overexpression seems to mediate many of the known phenotypic features associated with MYC, including promotion of apoptosis, alteration of morphology, enhancement of anchorage-independent growth, tumorigenic conversion, promotion of genomic instability, and inhibition of hematopoietic differentiation (By similarity).
------	--

基因ID	80177
基因名	MYCT1
Swiss	Q8N699
别名	MYCT1;MTLC

## 产品验证



Western blot analysis of MYCT1 expressed in Rat spleen, Rat lung using MYCT1 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)