

— ABLYBIO, Help Your Research



# HTATSF1 Rabbit pAb

货号: **AYP18914**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC IF/ICC
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IHC:</b> 1:50 - 1:100 <b>IF/ICC:</b> 1:50 - 1:200
理论分子量	85kDa
实测分子量	130kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,Jurkat,HepG2
细胞定位	Nucleus
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-180 of human HTATSF 1 (NP_055315.2).
------	--

## 靶点信息

研究背景	The protein encoded by this gene functions as a cofactor for the stimulation of transcriptional elongation by HIV-1 Tat, which binds to the HIV-1 promoter through Tat-TAR interaction. This protein may also serve as a dual-function factor to couple transcription and splicing and to facilitate their reciprocal activation. Alternatively spliced transcript variants have been found for this gene.
基因ID	27336
基因名	HTATSF1
Swiss	O43719 ( <a href="https://www.uniprot.org/uniprotkb/O43719/entry">https://www.uniprot.org/uniprotkb/O43719/entry</a> )
别名	HTATSF1,TAT-SF1,TATSF1,dj196E23.2,HTATSF1 Rabbit pAb,HIV Tat-specific factor 1

## 产品验证



Western blot analysis of HTATSF1 expressed in HeLa, Jurkat, HepG2 using HTATSF1 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn) (<https://www.ablybio.cn/>[www.ablybio.cn](http://www.ablybio.cn))