

ACOT2 Rabbit pAb

货号: **AYP18966**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200
理论分子量	32kDa/53kDa
实测分子量	50kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T,LO2,U-87MG,Mouse kidney
细胞定位	Mitochondrion
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 222-483 of human ACOT2 (NP_006812.3).
序列	IVDMFGTGGGLLEYRASLLAGKGFVVMALAYNYEDLPKTMETLHLEYFEEAMNYLLSHPEVKPGVGLLGISKGGELCLS MASFLKGITAAVVINGSVANVGGTLHYKGETLPPVGVNRRNIKVKTDGYADIVDVLNSPLEGPDQKSFIPVERAESTFLFLV GQDDHNWKSEFYANEACKRLQAHGRRKPKQIICYPETGHYIEPPYFPLCRASLHALVGSPHIIWGGEPRAHAMAQVDAWKQL QTFHFKHLGGHEGTIPSKV

靶点信息

研究背景	This gene encodes a member of the acyl-CoA thioesterase protein family, and is one of four acyl-CoA hydrolase genes located in a cluster on chromosome 14. Alternative splicing results in multiple transcript variants.
------	--

基因ID	10965
基因名	ACOT2
Swiss	P49753
别名	ACOT2;CTE-IA;CTE1A;MTE1;PTE2;PTE2A;ZAP128

产品验证



Western blot analysis of ACOT2 expressed in Mouse kidney,293T,LO2 using ACOT2 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn