

TRIM2 Rabbit pAb

货号: **AYP18969**

产品信息

反应	Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200
理论分子量	81kDa/84kDa
实测分子量	81kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse small intestine
细胞定位	Cytoplasm
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 100-300 of human TRIM 2 (NP_056086.2).
序列	GVAALQNNFFITNLMQVLRTPGSNAEESILETVTAVAAGKPLSCLPNDGDMVMEFYCQSCETAMCRECTEGEHAEHPTV PLKDVVEQHKASLQVQLDAVNKRLPEIDSALQFISEIHLQTLNQKASIVDDIHSTFDELQKTLNVRKSVLLMELEVNYGLKHK VLQSQLDTLLQGQESIKSCSNFTAQALNHGTETEVLLV

靶点信息

研究背景	The protein encoded by this gene is a member of the tripartite motif (TRIM) family. The TRIM motif includes three zinc-binding domains, a RING, a B-box type 1 and a B-box type 2, and a coiled-coil region. The protein localizes to cytoplasmic filaments. It plays a neuroprotective role and functions as an E3-ubiquitin ligase in proteasome-mediated degradation of target proteins. Mutations in this gene can cause early-onset axonal neuropathy. Alternative splicing results in multiple transcript variants.
------	---

基因ID	23321
基因名	TRIM2
Swiss	Q9C040
别名	TRIM2;CMT2R;RNF86

产品验证



Western blot analysis of TRIM2 expressed in Mouse small intestine using TRIM2 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn