

— ABLYBIO, Help Your Research



# PDIA5 Rabbit pAb

货号: **AYP18972**

## 产品信息

反应	Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	<b>WB:</b> 1:500 - 1:2000
理论分子量	30kDa/59kDa
实测分子量	59kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse kidney,Mouse liver,Mouse pancreas,Rat liver
细胞定位	Endoplasmic reticulum lumen
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 22-140 of human PDIA5 (NP_006801.1).
------	--

## 靶点信息

研究背景	This gene encodes a member of the disulfide isomerase (PDI) family of endoplasmic reticulum (ER) proteins that catalyze protein folding and thiol-disulfide interchange reactions. The encoded protein has an N-terminal ER-signal sequence, three catalytically active thioredoxin (TRX) domains, a TRX-like domain, and a C-terminal ER-retention sequence. The N-terminal TRX-like domain is the primary binding site for the major ER chaperone calreticulin and possibly other proteins and substrates as well. Alternative splicing results in multiple protein- and non-protein-coding transcript variants.
基因ID	10954
基因名	PDIA5
Swiss	Q14554 ( <a href="https://www.uniprot.org/uniprotkb/Q14554/entry">https://www.uniprot.org/uniprotkb/Q14554/entry</a> )
别名	PDIA5,PDIR,PDIA5 Rabbit pAb,Protein disulfide isomerase-related protein

## 产品验证



Western blot analysis of PDIA5 expressed in Mouse kidney,Rat liver,Mouse liver using PDIA5 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn) (<https://www.ablybio.cn/www.ablybio.cn>)