

— ABLYBIO, Help Your Research



IL17RE Rabbit pAb

货号: **AYP18999**

产品信息

| | |
|-------|------------------------------------------------------------------------------------------------------|
| 反应 | Human,Mouse |
| 宿主 | Rabbit |
| 克隆性 | Polyclonal |
| 预测反应 | |
| 应用 | WB |
| 推荐浓度 | WB: 1:500 - 1:2000 |
| 理论分子量 | 30kDa/43kDa/58kDa/61kDa/74kDa |
| 实测分子量 | 75kDa |
| 形式 | Liquid |
| 保存条件 | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3. |
| 偶联物 | Unconjugated |
| 阳性对照 | HT-29,A-549,A-431,Mouse kidney,Mouse thymus |
| 细胞定位 | Cell membrane,Cytoplasm,Secreted,Single-pass type I membrane protein |
| 纯化 | Affinity purification |


抗原信息

| | |
|------|----------------------------------------------------------------------------------------------------------------------|
| 抗原信息 | Recombinant fusion protein containing a sequence corresponding to amino acids 24-260 of human IL17R E (NP_705613.1). |
|------|----------------------------------------------------------------------------------------------------------------------|

靶点信息

| | |
|-------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| 研究背景 | This gene encodes a transmembrane protein that functions as the receptor for interleukin-17C. The encoded protein signals to downstream components of the mitogen activated protein kinase (MAPK) pathway. Activity of this protein is important in the immune response to bacterial pathogens. Alternatively spliced transcript variants have been described for this gene. |
| 基因ID | 132014 |
| 基因名 | IL17RE |
| Swiss | Q8NFR9 (https://www.uniprot.org/uniprotkb/Q8NFR9/entry) |
| 别名 | IL17RE,IL17RE Rabbit pAb |

产品验证



Western blot analysis of IL17RE expressed in Mouse kidney, HT-29, A-549 using IL17RE Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/>www.ablybio.cn)