

IL17RE Rabbit pAb

货号: **AYP18999**

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	30kDa/43kDa/58kDa/61kDa/74kDa
实测分子量	75kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HT-29,A-549,A-431,Mouse kidney,Mouse thymus
细胞定位	Cell membrane,Cytoplasm,Secreted,Single-pass type I membrane protein
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 24-260 of human IL17R E (NP_705613.1).
序列	AGIGFRHLPHWNTRCPLASHTDDSFSGSSAYIPCRTWWALFSTKPWCVRVWHCSRCLCQHLLSGGSLQRLGFHLLVQ KSKKSSTFKFYRRHKMPAPAQRKLLPRRHLSEKSHHISIPSPDISHKGLRSKRTQPSDPETWESLPRLDSQRHGGPEFSFD LLPEARAIRVTISSGPEVSVRLCHQWALECEELSSPYDVQKIVSGGHTVELPYEFLLPCLCIEASYLQEDTVRRKKCP

靶点信息

研究背景	This gene encodes a transmembrane protein that functions as the receptor for interleukin-17C. The encoded protein signals to downstream components of the mitogen activated protein kinase (MAPK) pathway. Activity of this protein is important in the immune response to bacterial pathogens. Alternatively spliced transcript variants have been described for this gene.
基因ID	132014

基因名	IL17RE
Swiss	Q8NFR9
别名	IL17RE

产品验证



Western blot analysis of IL17RE expressed in Mouse kidney, HT-29, A-549 using IL17RE Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn