

— ABLYBIO, Help Your Research



TNFRSF13C Rabbit pAb

货号: **AYP19008**

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	18kDa
实测分子量	38KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Raji,Daudi,SU-DHL-1
细胞定位	Membrane,Single-pass type III membrane protein
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-80 of human TNFRSF13C (NP_443177.1).
------	--

靶点信息

研究背景	B cell-activating factor (BAFF) enhances B-cell survival in vitro and is a regulator of the peripheral B-cell population. Overexpression of Baff in mice results in mature B-cell hyperplasia and symptoms of systemic lupus erythematosus (SLE). Also, some SLE patients have increased levels of BAFF in serum. Therefore, it has been proposed that abnormally high levels of BAFF may contribute to the pathogenesis of autoimmune diseases by enhancing the survival of autoreactive B cells. The protein encoded by this gene is a receptor for BAFF and is a type III transmembrane protein containing a single extracellular cysteine-rich domain. It is thought that this receptor is the principal receptor required for BAFF-mediated mature B-cell survival.
基因ID	115650
基因名	TNFRSF13C
Swiss	Q96RJ3 (https://www.uniprot.org/uniprotkb/Q96RJ3/entry)
别名	TNFRSF13C,BAFF-R,BAFFR,BROMIX,CD268,CVID4,prolixin,TNFRSF13C Rabbit pAb,B-cell-activating factor receptor,BAFF receptor,BLyS receptor 3,BR3

产品验证



Western blot analysis of TNFRSF13C expressed in Raji,Daudi,SU-DHL-1 using TNFRSF13C Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)