

— ABLYBIO, Help Your Research



IQWD1 Rabbit pAb

货号: **AYP19010**

产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	96kDa/98kDa/103kDa/106kDa
实测分子量	105kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal, 50% glycerol, pH7.3.
偶联物	Unconjugated
阳性对照	HeLa, LO2
细胞定位	Nucleus
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 460-650 of human IQWD1 (NP_001017977.1).
------	--

靶点信息

研究背景	The protein encoded by this gene is a ligand-dependent coactivator of nuclear receptors, including nuclear receptor subfamily 3 group C member 1 (NR3C1), glucocorticoid receptor (GR), and androgen receptor (AR). The encoded protein and DNA damage binding protein 2 (DDB2) may act as tumor promoters and tumor suppressors, respectively, by regulating the level of androgen receptor in prostate tissues. In addition, this protein can act with glucocorticoid receptor to promote human papillomavirus gene expression.
基因ID	55827
基因名	DCAF6
Swiss	Q58WW2 (https://www.uniprot.org/uniprotkb/Q58WW2/entry)
别名	DCAF6,1200006M05Rik,ARCAP,IQWD1,MSTP055,NRIP,PC326,IQWD1 Rabbit pAb,Androgen receptor complex-associated protein,IQ motif and WD repeat-containing protein 1,Nuclear receptor interaction protein

产品验证



Western blot analysis of IQWD1 expressed in HeLa, LO2 using IQWD1 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)