

SLCO3A1 Rabbit pAb

货号: **AYP19013**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	46kDa/62kDa/74kDa/76kDa
实测分子量	76kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	A-549,Mouse kidney,Mouse thymus,Mouse lung,Rat brain,Rat thymus
细胞定位	Cell membrane,Multi-pass membrane protein
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 440-540 of human SLCO3A1 (NP_037404.2).
序列	FLGCDTGPVAGVTVPYGNSTAPGSALDPYSPCINNCECQTDSTFVCGADGITYLSACFAGCNSTNLTGACLTTPAENATVWPGKCPSPGCQEAFITFL

靶点信息

研究背景	Mediates the Na(+)-independent transport of organic anions such as estrone-3-sulfate. Mediates transport of prostaglandins (PG E1 and E2, thyroxine (T4), deltorphin II, BQ-123 and vasopressin, but not DPDPE (a derivative of enkephalin lacking an N-terminal tyrosine residue, estrone-3-sulfate, taurocholate, digoxin or DHEAS).
基因ID	28232

基因名	SLCO3A1
Swiss	Q9UIG8
别名	SLCO3A1;OATP-D;OATP-RP3;OATP3A1;OATPD;OATPRP3;SLC21A11

产品验证



Western blot analysis of SLCO3A1 expressed in Mouse kidney,Rat brain,A-549 using SLCO3A1 Rabbit p Ab at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 12 0s.

实验步骤

访问官网浏览详情: www.ablybio.cn