

MLLT10 Rabbit pAb

货号: **AYP19027**

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	13kDa/20kDa/109kDa/113kDa
实测分子量	113kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T,Mouse spleen
细胞定位	Nucleus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 250-500 of human MLLT 10 (NP_001182555.1).
序列	EPSPALVPSLTVTTEKTYTSTSNNSISGSLKRLEDTTARFTNANFQEVS AHTSSGKDVSETRGSEGKGGKSSAHSSGQRG RKPGGGRNPGTTVSAASPFPQGSFSGTPGSVKSSSGSSVQSPQDFLSFTDSDLRNDYSYSHSQSSATKDVHKGESGSQ EGGVNSFSTLIGLPSTSAVTSQPKSFENSPGDLGNSLPTAGYKRAQTSGIEETVKEKKRKGKQSKHGPRPKGNKNQ ENVSHLSVSSASP

靶点信息

研究背景	This gene encodes a transcription factor and has been identified as a partner gene involved in several chromosomal rearrangements resulting in various leukemias. Multiple transcript variants encoding different isoforms have been found for this gene.
基因ID	8028

基因名	MLLT10
Swiss	P55197
别名	MLLT10;AF10

产品验证



Western blot analysis of MLLT10 expressed in 293T, Mouse spleen using MLLT10 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn