

HRH1 Rabbit pAb

货号: **AYP19042**

产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200
理论分子量	55kDa
实测分子量	56kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal, 50% glycerol, pH7.3.
偶联物	Unconjugated
阳性对照	Human normal colon tissue
细胞定位	Cell membrane, Multi-pass membrane protein
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 250-340 of human HRH 1 (NP_000852.1).
序列	KPGKESPWEVLKRKPKDAGGGSVLKSPSQTPKEMKSPVVFSDREDDREVDKLYCFPLDIVHMQAAAEGSSRDYVAVNRS HGQLKTDEQGLNT

靶点信息

研究背景	Histamine is a ubiquitous messenger molecule released from mast cells, enterochromaffin-like cells, and neurons. Its various actions are mediated by histamine receptors H1, H2, H3 and H4. The protein encoded by this gene is an integral membrane protein and belongs to the G protein-coupled receptor superfamily. It mediates the contraction of smooth muscles, the increase in capillary permeability due to contraction of terminal venules, the release of catecholamine from adrenal medulla, and neurotransmission in the central nervous system. It has been associated with multiple processes, including memory and learning, circadian rhythm, and thermoregulation. It is also known to contribute to the pathophysiology of allergic diseases such as atopic dermatitis, asthma, anaphylaxis and allergic rhinitis. Multiple alternatively spliced variants, encoding the same protein, have been identified.
基因ID	3269
基因名	HRH1
Swiss	P35367
别名	HRH1;H1-R;H1R;HH1R;hisH1

产品验证



Western blot analysis of HRH1 expressed in Human normal colon tissue using HRH1 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn