

— ABLYBIO, Help Your Research



CYP3A4 Rabbit pAb

货号: **AYP19046**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200
理论分子量	57kDa
实测分子量	57kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	rat liver tissue, mouse liver tissue, mouse brain tissue
细胞定位	Endoplasmic reticulum membrane, Microsome membrane, Single-pass membrane protein
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 244-503 of human CYP3A4 (NP_059488.2).
------	--

靶点信息

研究背景	This gene encodes a member of the cytochrome P450 superfamily of enzymes. The cytochrome P450 proteins are monooxygenases that catalyze many reactions involved in drug metabolism and synthesis of cholesterol, steroids and other lipids. This protein localizes to the endoplasmic reticulum and its expression is induced by glucocorticoids and some pharmacological agents. This enzyme is involved in the metabolism of approximately half the drugs in use today, including acetaminophen, codeine, cyclosporin A, diazepam and erythromycin. The enzyme also metabolizes some steroids and carcinogens. This gene is part of a cluster of cytochrome P450 genes on chromosome 7q21.1. Previously another CYP3A gene, CYP3A3, was thought to exist; however, it is now thought that this sequence represents a transcript variant of CYP3A4. Alternatively spliced transcript variants encoding different isoforms have been identified.
基因ID	1576
基因名	CYP3A4
Swiss	P08684 (https://www.uniprot.org/uniprotkb/P08684/entry)
别名	CYP3A4,CP33,CP34,CYP3A,CYP3A3,CYP3A3,CYP3A3,CYP3A3,CYP3A3,HLP,NF-25,P450C3,P450PCN1,CYP3A4 Rabbit pAb,1,4-cineole 2-exo-monooxygenase,8-cineole 2-exo-monooxygenase,Albendazole monooxygenase (sulfoxide-forming),Albendazole sulfoxidase,Cholesterol 25-hydroxylase,Cytochrome P450 3A3,Cytochrome P450 H Lp

产品验证



Western blot analysis of CYP3A4 expressed in rat liver tissue, mouse liver tissue, mouse brain tissue using CYP3A4 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/>www.ablybio.cn)