

PLCXD2 Rabbit pAb

货号: **AYP19056**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:100
理论分子量	34kDa
实测分子量	34kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	
细胞定位	nucleus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-305 of human PLCXD2 (NP_001172035.1).
序列	MLAVRKARRKLRMGTCSPNPSGKTTSSEVCNADWMASLPPHLHNLPLSNLAIPGSHDSFSYWVDEKSPVGPDPQTQAIKR LARISLVKMKLKKWSVTQNLTFREQLEAGIRYFDLRVSSKPGDADQEIYFIHGLFGIKVWDGLMEIDSFLTQHPQEIIFLDFN HFYAMDETHHKCLVLRIQEAFGNLCPACSVESLTLRTLWEKNCQVLIFYHCPFYKQYPFLWPGKKIPAWANTTSVRKLILF LETTLSERASRGSFHVSQAILTPRVKTIARGLVGGKNTLVHSNRWNSHGPSLLSQERS

靶点信息

研究背景	Predicted to enable phosphoric diester hydrolase activity. Predicted to be involved in lipid catabolic process and signal transduction. Predicted to be located in nucleus.
基因ID	257068

基因名	PLCXD2
Swiss	Q0VAA5
别名	PLCXD2

产品验证



Western blot analysis of PLCXD2 expressed in HUVEC, mouse brain, rat brain using PLCXD2 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn