

GALNT3 Rabbit pAb

货号: **AYP19109**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC IF/ICC
推荐浓度	WB: 1:500 - 1:1000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200
理论分子量	22kDa/72kDa
实测分子量	73KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	BxPC-3,Rat testis
细胞定位	Golgi apparatus,Golgi stack membrane,Single-pass type II membrane protein
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-140 of human GALNT3 (NP_004473.2).
序列	MAHLKRLVKLHIKRHYHKKFWKLGAVIFFFIIVLVMQREVSQYSKEESRMERNMKNKMKMLDLMLEAVNNIKDAMPKM QIGAPVRQNIDAGERPCLQGYTAAELKPVLDLPPQDSNAPGASGKAFKTTNLSVEEQKE

靶点信息

研究背景	This gene encodes UDP-GalNAc transferase 3, a member of the GalNAc-transferases family. This family transfers an N-acetyl galactosamine to the hydroxyl group of a serine or threonine residue in the first step of O-linked oligosaccharide biosynthesis. Individual GalNAc-transferases have distinct activities and initiation of O-glycosylation is regulated by a repertoire of GalNAc-transferases. The protein encoded by this gene is highly homologous to other family members, however the enzymes have different substrate specificities.
------	--

基因ID	2591
基因名	GALNT3
Swiss	Q14435
别名	GALNT3;GalNAc-T3;HFTC;HHS

产品验证



Western blot analysis of GALNT3 expressed in BxPC-3, Rat testis using GALNT3 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn