

— ABLYBIO, Help Your Research



FLJ12529 Rabbit pAb

货号: **AYP19151**

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	51kDa/52kDa/56kDa
实测分子量	60kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Jurkat,HepG2,Mouse thymus
细胞定位	Nucleus
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 50-240 of human FLJ12529 (NP_079087.3).
------	---

靶点信息

研究背景	Cleavage factor Im (CFIm) is one of six factors necessary for correct cleavage and polyadenylation of pre-mRNAs. CFIm is composed of three different subunits of 25, 59, and 68 kDa, and it functions as a heterotrimer, with a dimer of the 25 kDa subunit binding to two of the 59 or 68 kDa subunits. The protein encoded by this gene represents the 59 kDa subunit, which can interact with the splicing factor U2 snRNP Auxiliary Factor (U2AF) 65 to link the splicing and polyadenylation complexes.
基因ID	79869
基因名	CPSF7
Swiss	Q8N684 (https://www.uniprot.org/uniprotkb/Q8N684/entry)
别名	CPSF7,CFIm59,FLJ12529,FLJ12529 Rabbit pAb,Cleavage and polyadenylation specificity factor 59 kDa subunit,Cleavage factor Im complex 59 kDa subunit,Pre-mRNA cleavage factor Im 59 kDa subunit

产品验证



Western blot analysis of FLJ12529 expressed in Jurkat, HepG2, Mouse thymus using FLJ12529 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/>www.ablybio.cn)