

— ABLYBIO, Help Your Research



DLGAP2 Rabbit pAb

货号: **AYP19181**

产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	108kDa/115kDa/117kDa
实测分子量	117kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal, 50% glycerol, pH7.3.
偶联物	Unconjugated
阳性对照	U-87MG, 293T
细胞定位	Cell junction, Cell membrane, Peripheral membrane protein, postsynaptic cell membrane, postsynaptic density, synapse
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 550-800 of human DLGAP2 (NP_004736.2).
------	--

靶点信息

研究背景	The product of this gene is a membrane-associated protein that may play a role in synapse organization and signalling in neuronal cells. This gene is biallelically expressed in the brain, however, only the paternal allele is expressed in the testis (PMID:18055845). Alternatively spliced transcript variants encoding different isoforms have been identified.
基因ID	9228
基因名	DLGAP2
Swiss	Q9P1A6 (https://www.uniprot.org/uniprotkb/Q9P1A6/entry)
别名	DLGAP2,C8orf68,DAP2,ERICH1-AS1,SAPAP2,DLGAP2 Rabbit pAb,PSD-95/SAP90-binding protein 2,SAP90/PSD-95-associated protein 2

产品验证



Western blot analysis of DLGAP2 expressed in U-87MG,293T using DLGAP2 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)