

— ABLYBIO, Help Your Research



FCRL5 Rabbit pAb

货号: **AYP19185**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	14kDa/17kDa/65kDa/83kDa/106kDa
实测分子量	106kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Jurkat,PC-12,SP2/0
细胞定位	Cell membrane,Single-pass type I membrane protein
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 16-190 of human FCRL5 (NP_001182317.1).
------	---

靶点信息

研究背景	This gene encodes a member of the immunoglobulin receptor superfamily and the Fc-receptor like family. This gene and several other Fc receptor-like gene members are clustered on the long arm of chromosome 1. The encoded protein is a single-pass type I membrane protein and contains 8 immunoglobulin-like C2-type domains. This gene is implicated in B cell development and lymphomagenesis. Alternatively spliced transcript variants encoding different isoforms have been identified.
基因ID	83416
基因名	FCRL5
Swiss	Q96RD9 (https://www.uniprot.org/uniprotkb/Q96RD9/entry)
别名	FCRL5,BXMAS1,CD307,CD307e,FCRH5,IRTA2,PRO820,FCRL5 Rabbit pAb,Fc receptor homolog 5,Immune receptor translocation-associated protein 2

产品验证



Western blot analysis of FCRL5 expressed in Jurkat,PC-12,SP2/0 using FCRL5 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn (<https://www.ablybio.cn/www.ablybio.cn>)