

IFIT2 Rabbit pAb

货号: **AYP19203**

产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	54kDa
实测分子量	50kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal, 50% glycerol, pH7.3.
偶联物	Unconjugated
阳性对照	A-431, HepG2
细胞定位	Cytoplasm, Endoplasmic reticulum
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 273-472 of human IFIT2 (NP_001538.4).
序列	ALEYIPNNAYLHCQIGCCYRAKVFQVMNLRNGMYGKRKLELIGHAVAHLKKADEANDNLFVCSILASLHALADQYEDAE YYFQKEFSKELTPVAKQLLHLRYGNFQLYQMKCEDKAIHHFIEGVKINQKSREKEKMKDKLQKIAKMRLSKNGADSEALHV LAFLQELNEKMQQAEDSERGLESGSLIPSASSWNGE

靶点信息

研究背景	IFN-induced antiviral protein which inhibits expression of viral messenger RNAs lacking 2'-O-methylation of the 5' cap. The ribose 2'-O-methylation would provide a molecular signature to distinguish between self and non-self mRNAs by the host during viral infection. Viruses evolved several ways to evade this restriction system such as encoding their own 2'-O-methylase for their mRNAs or by stealing host cap containing the 2'-O-methylation (cap snatching mechanism). Binds AU-rich viral RNAs, with or without 5' triphosphorylation, RNA-binding is required for antiviral activity. Can promote apoptosis.
------	---

基因ID	3433
基因名	IFIT2
Swiss	P09913
别名	IFIT2;G10P2;GARG-39;IFI-54;IFI-54K;IFI54;IFIT-2;ISG-54 K;ISG-54K;ISG54;P54;cig42

产品验证



Western blot analysis of IFIT2 expressed in A-431, HepG2 using IFIT2 Rabbit pAb at 1:1000. Secondary anti body: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% no n-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn