

SLC25A14 Rabbit pAb

货号: **AYP19209**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200
理论分子量	35kDa/36kDa/39kDa
实测分子量	36kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse brain,Rat brain
细胞定位	Mitochondrion inner membrane,Multi-pass membrane protein
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 12-262 of human SLC25A14 (NP_001269127.1).
序列	GGLASIVAEFGTFFPVDLTKTRLQVQGQSIDARFKEIKYRGMFHALFRICKEEGLVLYSGIAPALLRQASYGTIKIGIYQSLKRLFVERLEDETLLINMICGVVSGVISSTIANPTDVLKIRMQAQGSLSFQGSIMIGSFIDIYQQEGTRGLWRGVVPTAQRAAIVGV VELPVYDITKKHLILSGMMGDTILTHFVSSFTCGLAGALASNPVDVVRTRMMNQRAIVGHVDLYKGTVDGILKMWKHEGF FALYK

靶点信息

研究背景	Mitochondrial uncoupling proteins (UCP) are members of the larger family of mitochondrial anion carrier proteins (MACP). Uncoupling proteins separate oxidative phosphorylation from ATP synthesis with energy dissipated as heat, also referred to as the mitochondrial proton leak. Uncoupling proteins facilitate the transfer of anions from the inner to the outer mitochondrial membrane and the return transfer of protons from the outer to the inner mitochondrial membrane. They also reduce the mitochondrial membrane potential in mammalian cells. This gene is widely expressed in many tissues with the greatest abundance in brain and testis. Alternative splicing results in multiple transcript variants. A pseudogene of this gene has been defined on chromosome 4.
基因ID	9016
基因名	SLC25A14
Swiss	O95258
别名	SLC25A14;BMCP1;UCP5

产品验证



Western blot analysis of SLC25A14 expressed in Mouse brain,Rat brain using SLC25A14 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn