

TNFAIP8L2 Rabbit pAb

货号: **AYP19228**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IF/ICC IP
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200 IP: 1:500 - 1:1000
理论分子量	20kDa
实测分子量	18KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	THP-1,Mouse thymus,Mouse spleen,Rat spleen
细胞定位	cytoplasm
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-184 of human TNFAIP8L2 (NP_078851.2).
序列	MESFSSKSLALQAEKKLLSKMAGRSVAHLFIDETSSEVLDELRYVSKEYTHSRPQAQRVIKDLIKVAIKVAVLHRNGSFGPS ELALATRFQRQKLRQGAMTALSFGVEVDFTEAAVLAGLLTECRDVLLLEVEHHLTPKSHGRIRHVDFHSDPGLLTALYGPDF TQHLGKICDGLRKLLEDEGKL

靶点信息

研究背景	Acts as a negative regulator of innate and adaptive immunity by maintaining immune homeostasis. Negative regulator of Toll-like receptor and T-cell receptor function. Prevents hyperresponsiveness of the immune system and maintains immune homeostasis. Inhibits JUN/AP1 and NF-kappa-B activation. Promotes Fas-induced apoptosis (By similarity).
------	--

基因ID	79626
基因名	TNFAIP8L2
Swiss	Q6P589
别名	TNFAIP8L2;TIPE2

产品验证



Western blot analysis of TNFAIP8L2 expressed in Mouse thymus, Rat spleen, THP-1 using TNFAIP8L2 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn