

# Uhrf2 Rabbit pAb

货号: **AYP19250**

## 产品信息

反应	Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	<b>WB:</b> 1:500 - 1:2000
理论分子量	24kDa/57kDa/90kDa
实测分子量	90kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse thymus,Mouse testis,Mouse liver,Rat thymus
细胞定位	Nucleus
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 76-359 of mouse Uhrf2 ( NP_659122.2).
序列	DSSLPSTSKQNDAQVKPSSHPPKVKKTARGGSSSQPSTSARTCLIDPGFGLYKVNELVDARDVGLGAWFEAHIHSVTR ASDGH5RGTPLKNGSSYKRTNGNVNHNSKENTNKLDNVPSTSN5SDSVAADEDVIYHIEYDEYPESGILEMNVKDLRPRA RTILKWNELNVGDVVMVNYNVENPGKRGFWYDAEITTLKTISRKKKEVRVKVFLGGSEGTLNDCRVMSVDEIFKIEKPGAH PISFADGKFLRKN2PECDLCCGGDPDKTCHMCSCHKCGEKRPNM

## 靶点信息

研究背景	This gene encodes a nuclear protein which is involved in cell-cycle regulation. The encoded protein is a ubiquitin-ligase capable of ubiquinating PCNP (PEST-containing nuclear protein), and together they may play a role in tumorigenesis. The encoded protein contains an NIRF_N domain, a PHD finger, a set- and ring-associated (SRA) domain, and a RING finger domain and several of these domains have been shown to be essential for the regulation of cell proliferation. This protein may also have a role in intranuclear degradation of polyglutamine aggregates. Alternative splicing results in multiple transcript variants some of which are non-protein coding.
基因ID	109113
基因名	
Swiss	Q7TMI3
别名	UHRF2;NIRF;RNF107;TDRD23;URF2;Uhrf2

## 产品验证



Western blot analysis of Uhrf2 expressed in Mouse thymus, Rat thymus, Mouse testis using Uhrf2 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120 S.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)