

PROCR Rabbit pAb

货号: **AYP19256**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC
推荐浓度	WB: 1:500 - 1:1000 IHC: 1:50 - 1:200
理论分子量	26kDa
实测分子量	45KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	U-87MG
细胞定位	Membrane,Single-pass type I membrane protein
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 19-209 of human PROC R (NP_006395.2).
序列	QDASDGLQRLHMLQISYFRDPYHVWYQGNASLGHLTHVLEGPDTNTTIIQLQPLQEPESWARTQSGLQSYLLQFHGLV RLVHQERTLAFPLTIRCFLGCELPPEGSRAHVFFEVAVNGSSFVSRPERALWQADTQVTSQVVTFTLQQLNAYNRTRYEL REFLEDTCVQYVQKHISAENTKGSQTSRSYT

靶点信息

研究背景	The protein encoded by this gene is a receptor for activated protein C, a serine protease activated by and involved in the blood coagulation pathway. The encoded protein is an N-glycosylated type I membrane protein that enhances the activation of protein C. Mutations in this gene have been associated with venous thromboembolism and myocardial infarction, as well as with late fetal loss during pregnancy. The encoded protein may also play a role in malarial infection and has been associated with cancer.
------	--

基因ID	10544
基因名	PROCR
Swiss	Q9UNN8
别名	PROCR;CCCA;CCD41;EPCR;h-PROCR

产品验证



Western blot analysis of PROCR expressed in U-87MG using PROCR Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn