

# CXXC1 Rabbit pAb

货号: AYP19330

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IHC:</b> 1:50 - 1:200
理论分子量	75kDa/76kDa
实测分子量	76kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HepG2
细胞定位	Nucleus speckle
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-210 of human CXXC1 (NP_001095124.1).
序列	MEGDGSDPEPPDAGEDSKSENGENAPIYCICRKPDPDINCFMIGCDNCNEWFHGDCIRITEKMAKAIREWYCRECREKDPKLE IRYRHKKSRERDGNERDSSEPRDEGGGRKRPVDPDLQRRAGSGTGVGAMLARGSASPHKSSPQLVATPSQHHQQQ QQQIKRSARMCGECEACRRTEDCGHCDFCRDMKKFGGPNKIRQKCLRQCQL

## 靶点信息

研究背景	This gene encodes a protein that functions as a transcriptional activator that binds specifically to non-methylated CpG motifs through its CXXC domain. The protein is a component of the SETD1 complex, regulates gene expression and is essential for vertebrate development.
基因ID	30827

基因名	CXXC1
Swiss	Q9P0U4
别名	CXXC1;2410002I16Rik;5830420C16Rik;CFP1;CGBP;HsT2645;PCCX1;PHF18;SPP1;ZCGPC1;hCGBP

## 产品验证



Western blot analysis of CXXC1 expressed in HepG2 using CXXC1 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)