

POLR2H Rabbit pAb

货号: **AYP19366**

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IP
推荐浓度	WB: 1:500 - 1:2000 IP: 1:50 - 1:200
理论分子量	9kDa/12kDa/13kDa/17kDa/19kDa
实测分子量	17kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	SW480,MCF7,HL-60,293T,Mouse spleen,Mouse thymus
细胞定位	Nucleus,nucleolus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-150 of human POLR2 H (NP_006223.2).
序列	MAGILFEDIFDVKIDIDPEGKKFDRVSRRLHCESESFKMDLILDVNIQIYPVDLGDKFRLVIASSTLYEDGTLDGGEYNPTDDRP SRADQFEYVMYGKVYRIEGDETSTEAAATRLSAYVSYGGLLMRLQGDANNLHGFEVDSRVYLLMKKLAF

靶点信息

研究背景	The three eukaryotic RNA polymerases are complex multisubunit enzymes that play a central role in the transcription of nuclear genes. This gene encodes an essential and highly conserved subunit of RNA polymerase II that is shared by the other two eukaryotic DNA-directed RNA polymerases, I and III. Alternative splicing results in multiple transcript variants of this gene.
基因ID	5437

基因名	POLR2H
Swiss	P52434
别名	POLR2H;RPABC3;RPB17;RPB8

产品验证



Western blot analysis of POLR2H expressed in Mouse spleen, SW480, MCF7 using POLR2H Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn