

# THOC3 Rabbit pAb

货号: **AYP19403**

## 产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	<b>WB:</b> 1:500 - 1:2000
理论分子量	36kDa/38kDa
实测分子量	39kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	LO2,Mouse brain
细胞定位	Nucleus,Nucleus speckle
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-200 of human THOC3 (NP_115737.1).
序列	MAVPAAAMGPSALGQSGPGSMAPWCSVSSGSPRYVLGMQELFRGHSKTREFLAHSAKVHSAVWASCDGRRRLASGSFDK TASVFLLEKDRDLVKENNYRGHGSDVDQLCWHPSPDLFVTASGDKTIRIWDVVRTTKCIATVNTKGENINICWSPDGQTIAV GNKDDVVTFIDAKTHRSKAEEQFKFEVNEISWNNDNNMFFLT

## 靶点信息

研究背景	This gene encodes a component of the nuclear THO transcription elongation complex, which is part of the larger transcription export (TREX) complex that couples messenger RNA processing and export. In human s, the transcription export complex is recruited to the 5'-end of messenger RNAs in a splicing- and cap-dependent manner. Studies of a related complex in mouse suggest that the metazoan transcription export c complex is involved in cell differentiation and development. A pseudogene of this gene has been defined o n chromosome 5.
------	--

基因ID	84321
基因名	THOC3
Swiss	Q96J01
别名	THOC3;THO3;hTRESK45

## 产品验证



Western blot analysis of THOC3 expressed in LO2, Mouse brain using THOC3 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)