

RBM11 Rabbit pAb

货号: **AYP19409**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	26kDa/32kDa
实测分子量	32kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	U-87MG,293T,MCF7,Mouse testis,Rat testis
细胞定位	Nucleus,Nucleus speckle,nucleoplasm
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-281 of human RBM11 (NP_658983.3).
序列	MFPAQEEADRTVFGNLEARVREEILYELFLQAGPLTKVTICKDREGKPKSFGFVCFKHPESVSYAIALINGIRLYGRPINVQY RFGSSRSSEPANQSFESCCKINSHNYRNEEMLVGRSSFPQYFIPINNTSLPQEYFLFQKMQWHVYNPVLQLPYYEMTAPL PNSASVSSSLNHVPDLEAGPSSYKWITHQQPSDSDLYQMTAPLPNSASVSSSLNHVPDLEAGPSSYKWITHQQPSDSDLY QMNKRKRQKQTSDDSSTDNRRGNECSQKFRKSKKKKRY

靶点信息

研究背景	Tissue-specific splicing factor with potential implication in the regulation of alternative splicing during neuron and germ cell differentiation. Antagonizes SRSF1-mediated BCL-X splicing. May affect the choice of alternative 5' splice sites by binding to specific sequences in exons and antagonizing the SR protein SRSF1.
基因ID	54033

基因名	RBM11
Swiss	P57052
别名	RBM11

产品验证



Western blot analysis of RBM11 expressed in Mouse testis,Rat testis,U-87MG using RBM11 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn