

DCAF13 Rabbit pAb

货号: **AYP19413**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	10kDa/51kDa
实测分子量	51kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	PC-12,NIH/3T3
细胞定位	Nucleus,nucleolus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 298-597 of human DCA F13 (NP_056235.4).
序列	EEEPLHTILGKTVYTGIDHHWKEAVFATCGQQVDIWDEQRTNPICSMWTGFDSSISVKFNPIETFLGSCASDRNIVLYDM RQATPLKKVILDMRTNTICWNPMEAFIFTAANEDYNLYTFDMRALDTPVMVHMDHVSALVDVDYSPTGKEFVSASFDKSI RIFPVDKSRSEVYHTKRMQHVICVKWTSDSKYIMCGSDEMNIWLKANASEKLGVLTSREKAAKDYNQKLKEKFQHYPH IKRIARHRHLPKSIYSQIQEQRIMKEARRRKEVNRIRKHSKPGSVPLVSEKKKHVVAVVK

靶点信息

研究背景	Possible role in ribosomal RNA processing (By similarity. May function as a substrate receptor for CUL4-D DB1 E3 ubiquitin-protein ligase complex.
基因ID	25879

基因名	DCAF13
Swiss	Q9NV06
别名	DCAF13;GM83;HSPC064;WDSOF1;Sof1

产品验证



Western blot analysis of DCAF13 expressed in PC-12,NIH/3T3 using DCAF13 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn