

# NME1-NME2 Rabbit pAb

货号: **AYP19453**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	<b>WB:</b> 1:500 - 1:2000
理论分子量	22kDa
实测分子量	17kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	U-87MG,OVCAR3,Mouse kidney,Rat kidney,Rat liver
细胞定位	Cell projection,Cytoplasm,Nucleus,lamellipodium,ruffle
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-267 of human NME1-NME2 (NP_001018146.1).
序列	MANCERTFIAIKPDGVQRGLVGEIIRFEQKGFRLVGLKFMQASEDLLKEHYVDLKDRPFFAGLVKYMHS GPVVAMVWEG LNVVKTGRVMLGETNPADSKPGTIRGDFCIQVGRMTANLERTFIAIKPDGVQRGLVGEIIRFEQKGFRLVAMKFLRASEEH LKQHYIDLKDRPFFGLVKYMN SGPVVAMVWEG LNVVKTGRVMLGETNPADSKPGTIRGDFCIQVGRNIIHGSDSVKSAE KEISLWFKPEELVDYKSCAHDWVYE

## 靶点信息

研究背景	This locus represents naturally occurring read-through transcription between the neighboring NME1 and NME2 genes. The significance of this read-through transcription and the function of the resulting protein product have not yet been determined. Alternative splicing of this gene results in multiple transcript variants.
------	--

基因ID	654364
基因名	
Swiss	
别名	NME1-NME2;NM23-LV;NMELV

## 产品验证



Western blot analysis of NME1-NME2 expressed in Mouse kidney,Rat kidney,U-87MG using NME1-NME2 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/prot eins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)