

FMN1 Rabbit pAb

货号: **AYP19455**

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	53kDa/71kDa/132kDa/157kDa
实测分子量	158kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T,A-549,U-87MG,LO2,Mouse testis
细胞定位	Cell junction,Cell membrane,Cytoplasm,Cytoplasmic side,Nucleus,Peripheral membrane protein,adherens junction
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 350-495 of human FMN 1 (NP_001096654.1).
序列	NIDMPKTEPKGADPESPRREEMGCNADQESQSGPGVPQTQGGEVKPKSPETALEAFKALFIRPPRKGTTADTSELEALKR KMRHEKESLRAVFERSNSKPADGSPDSKSPDHSLTEQDDRTPGRLQAVWPPPKTKDTEEKVGLKYT

靶点信息

研究背景	This gene belongs to the formin homology family and encodes a protein that has a role in the formation of adherens junction and the polymerization of linear actin cables. The homologous gene in mouse is associated with limb deformity. Alternatively spliced transcript variants have been found for this gene.
基因ID	342184

基因名	FMN1
Swiss	Q68DA7
别名	FMN1;FMN;LD;formin-1

产品验证



Western blot analysis of FMN1 expressed in Mouse testis,293T,A-549 using FMN1 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn